arXiv:1704.04392v1 [math.FA] 14 Apr 2017

A NOTE ON TRIANGULAR OPERATORS ON SMOOTH SEQUENCE SPACES

ELIF UYANIK AND MURAT H. YURDAKUL

ABSTRACT. For a scalar sequence $(\theta_n)_{n\in\mathbb{N}}$, let C be the matrix defined by $c_n^k = \theta_{n-k+1}$ if $n \geq k$, $c_n^k = 0$ if $n < k$. The map between Köthe spaces $\lambda(A)$ and $\lambda(B)$ is called a Cauchy Product map if it is determined by the triangular matrix C. In this note we introduced some necessary and sufficient conditions for a Cauchy Product map on a nuclear Köthe space $\lambda(A)$ to nuclear G_1-space $\lambda(B)$ to be linear and continuous. Its transpose is also considered.

Dedicated to the memory of Prof. Dr. Tosun Terzioğlu

1. Introduction

We refer the reader to [3], [4] and [5] for the terminology used but not defined here. Let $A = (a_n^k)_{n,k\in\mathbb{N}}$ be a matrix of real numbers such that $0 \leq a_n^k \leq a_n^{k+1}$ for all n, k and $\sup_k a_n^k > 0$. The ℓ^1 - Köthe space $\lambda(A)$ defined by the matrix A is the space of all sequences of scalars $x = (x_n)$ such that

$$\|x\|_k = \sum_n |x_n| a_n^k < \infty, \quad \forall k \in \mathbb{N}$$

With the topology generated by the system of seminorms $\{\|.\|_k, k \in \mathbb{N}\}$, it is a Fréchet space.

The topological dual of $\lambda(A)$ is isomorphic to the space of all sequences u for which $|u_n| \leq C a_n^k$ for some k and $C > 0$.

It is well known that a Köthe space $\lambda(A)$ associated with the matrix A is nuclear if and only if for each k there exists m such that

$$\sum_n \frac{a_n^k}{a_n^m} < +\infty$$

2010 *Mathematics Subject Classification.* 47B37, 46A45.

Key words and phrases. Köthe spaces, Smooth Sequence Spaces, Cauchy Product.

This research was partially supported by the Turkish Scientific and Technological Research Council.

and in this case the fundamental system of norms $\|x\|_k = \sum_n |x_n| a_n^k$ can be replaced by the equivalent system of norms

$$\|x\|_k = \sup_n |x_n| a_n^k, \quad k \in \mathbb{N}.$$

The infinite and finite type power series spaces are well known examples of Köthe spaces given by the matrices $(e^{k\alpha_n})$ respectively $(e^{-\frac{\alpha_n}{k}})$ where (α_n) is a monotonically increasing sequence going to infinity. The space $A(\mathbb{C})$ of all entire functions on $\mathbb{C}$ and the space $A(\mathbb{D})$ of all holomorphic functions on the unit disc can be represented as an infinite respectively finite type power series spaces.

Smooth sequence spaces were introduced in [6] as a generalization of power series spaces. A Köthe set $A = \left\{(a_n^k)\right\}$ is called a G_∞-set and the corresponding Köthe space $\lambda(A)$ a G_∞-space if A satisfies the followings :

(1) $a_n^1 = 1$, $a_n^k \leq a_{n+1}^k$ for each k and n;

(2) $\forall k\ \exists j$ with $(a_n^k)^2 = O(a_n^j)$

A Köthe set $B = \left\{(b_n^k)\right\}$ is called a G_1-set and the corresponding Köthe space $\lambda(B)$ a G_1-space if B satisfies the followings :

(1) $0 < b_{n+1}^k \leq b_n^k < 1$ for each k and n;

(2) $\forall k\ \exists j$ with $b_n^k = O((b_n^j)^2)$

We need the following result [1].

Lemma 1.1. *Let $\lambda(A)$ and $\lambda(B)$ be Köthe spaces. A map $T : \lambda(A) \longrightarrow \lambda(B)$ is continuous linear map if and only if for each k there exists m such that*

$$\sup_n \frac{\|Te_n\|_k}{\|e_n\|_m} < +\infty$$

If (a_n), (b_n) are two sequences of scalars, then the Cauchy product $(c_n) = (a_n) * (b_n)$ of (a_n) and (b_n) is defined by $c_n = \sum_{k=1}^{n} a_{n+1-k} b_k$.

Now let $\theta = (\theta_n)$ be a fixed sequence of scalars and let $\lambda(A)$, $\lambda(B)$ be two nuclear ℓ^1-Köthe spaces. We define the Cauchy Product mapping T_θ from $\lambda(A)$ into $\lambda(B)$ by $T_\theta x = \theta * x, \quad x = (x_n) \in \lambda(A)$. So, $T_\theta : \lambda(A) \longrightarrow \lambda(B)$ can be determined by the lower triangular matrix

$$C = \begin{pmatrix} \theta_1 & 0 & 0 & 0 & \cdots \\ \theta_2 & \theta_1 & 0 & 0 & \cdots \\ \theta_3 & \theta_2 & \theta_1 & 0 & \cdots \\ \vdots & & & \ddots & \end{pmatrix}$$

2. Cauchy Product Map on Köthe spaces

In this section we introduce some necessary and sufficient conditions for the map T_θ to be linear and continuous.

Theorem 2.1. *Let $\lambda(A)$ be a nuclear Köthe space, $\lambda(B)$ be a nuclear G_1-space. Then the Cauchy product map $T_\theta : \lambda(A) \longrightarrow \lambda(B)$ is linear continuous operator if and only if the following hold:*
i) $\theta \in \lambda(B)$
ii) $\lambda(A) \subset \lambda(B)$

Proof. Let $T_\theta : \lambda(A) \longrightarrow \lambda(B)$ be a continuous linear operator. Note that $\|T_\theta e_n\|_k = \|(0, 0, ..., 0, \theta_1, \theta_2, \cdots)\|_k = \sup_{j\geq n} |\theta_{j-n+1}| b_j^k$ for $n \in \mathbb{N}$. Clearly $\|e_n\|_m = a_n^m$. So, by Lemma 1.1 $\forall k$, $\exists m$, $\exists \rho > 0$ such that

$$\sup_{j\geq n} |\theta_{j-n+1}| b_j^k \leq \rho a_n^m, \quad \forall n \in \mathbb{N}.$$

Choose $j = n$. Then $\forall k$, $\exists m$, $\exists C > 0$ such that

$$b_n^k \leq C a_n^m$$

i.e, $\lambda(A) \subset \lambda(B)$. Since $T_\theta e_1 \in \lambda(B)$, it follows that $\theta \in \lambda(B)$.

Conversely, since B is a G_1-set and by ii and i we have for a given k, there are $m_1(k)$ and $m_2(m_1)$ such that

$$\begin{aligned}
\|T_\theta e_n\|_k &= \sup_{j\geq n} |\theta_{j-n+1}| b_j^k \\
&\leq C_1 \sup_{j\geq n} |\theta_{j-n+1}| (b_j^{m_1})^2 \\
&\leq C_1 \sup_{j\geq n} (|\theta_{j-n+1}| b_j^{m_1})(b_n^{m_1}) \\
&\leq C_2 \sup_{j\geq n} (|\theta_{j-n+1}| b_j^{m_1})(a_n^{m_2}) \\
&\leq C_2 \sup_{j\geq n} (|\theta_{j-n+1}| b_{j-n+1}^{m_1})(a_n^{m_2}) \\
&\leq C a_n^{m_2}.
\end{aligned}$$

Therefore, $\forall k$, $\exists m_2$ such that

$$\sup_n \frac{\|T_\theta e_n\|_k}{\|e_n\|_{m_2}} < \infty$$

that is, T_θ is continuous. □ □

We consider the map $T_\theta' : \lambda(A) \longrightarrow \lambda(B)$ which is determined by the matrix C^t (the transpose of C) and try to find necessary and sufficient conditions for the continuity of T_θ'.

Theorem 2.2. *Let $\lambda(A)$ be a nuclear G_∞-space, $\lambda(B)$ be a nuclear Köthe space. Then, $T_\theta' : \lambda(A) \longrightarrow \lambda(B)$ which is given above is linear continuous operator if and only if the following hold:*
i) $\theta \in \lambda(A)'$
ii) $\lambda(A) \subset \lambda(B)$

Proof. The matrix C^t of the operator $T_\theta' : \lambda(A) \longrightarrow \lambda(B)$ is the following upper triangular matrix:

$$C^t = \begin{pmatrix} \theta_1 & \theta_2 & \theta_3 & \theta_4 & \cdots \\ 0 & \theta_1 & \theta_2 & \theta_3 & \cdots \\ 0 & 0 & \theta_1 & \theta_2 & \cdots \\ \vdots & & & \ddots & \end{pmatrix}$$

Let $T_\theta' : \lambda(A) \longrightarrow \lambda(B)$ be a continuous linear operator.
Note that $\|T_\theta' e_n\|_k = \|(\theta_n, \theta_{n-1}, \cdots, \theta_1, 0, 0, \cdots)\|_k = \sup_{1\leq i\leq n} |\theta_{n+1-i}| b_i^k$ for $n \in \mathbb{N}$. So, by Lemma 1.1 $\forall k$, $\exists m$, $\exists \mu > 0$ such that

$$\sup_{1\leq i\leq n} |\theta_{n+1-i}| b_i^k \leq \mu a_n^m, \quad \forall n \in \mathbb{N}.$$

Let $i = 1$. Hence $\exists m$, $\exists C = \frac{\mu}{b_1^k} > 0$ such that

$$|\theta_n| \leq C a_n^m, \quad \forall n$$

i.e, $\theta \in \lambda(A)'$.
Let $i = n$. Then $\forall k$, $\exists m$ such that

$$b_n^k \leq \frac{\mu}{|\theta_1|} a_n^m$$

i.e,

$$\lambda(A) \subset \lambda(B)$$

On the other hand, since A is a G_∞-set and by (i) and (ii) for a given k, there are m_1 and $m_2(k)$ and $m = max\{m_1, m_2\}$ such that

$$\begin{aligned} \|T_\theta' e_n\|_k &= \sup_{1\leq i\leq n} |\theta_{n-i+1}| b_i^k \\ &\leq C_1 \sup_{1\leq i\leq n} a_{n-i+1}^{m_1} b_i^k \\ &\leq C_1 \sup_{1\leq i\leq n} a_{n-i+1}^{m_1} a_i^{m_2} \\ &\leq C_1 a_n^{m_1} a_n^{m_2} \\ &\leq C_2 (a_n^m)^2. \end{aligned}$$

Since $\lambda(A)$ is G_∞ - space, for this m, $\exists j$ such that

$$\sup_n \frac{(a_n^m)^2}{a_n^j} < \infty$$

Therefore, $\forall k$, $\exists j$ such that

$$\sup_n \frac{\|T_\theta' e_n\|_k}{\|e_n\|_j} < \infty$$

that is, T_θ' is continuous. □ □

It is known that $\mathcal{S}$ is a normal sequence space if whenever $|x_i| < |y_i|$ and $y = (y_i) \in \mathcal{S}$, then $x = (x_i) \in \mathcal{S}$ [2].

Remark 2.1. *Now we write $\theta \in \mathcal{S}$ when the Cauchy product map $T_\theta : \lambda(A) \longrightarrow \lambda(B)$ above is continuous. If $\theta, \eta \in \mathcal{S}$, $\lambda \in \mathcal{K}$, then clearly $T_{\theta+\eta}$ and $T_{\lambda\theta}$ will be continuous since T_θ and T_η are continuous. Hence $\mathcal{S}$ is a vector space.*

Now, let $|\theta_i| < |\eta_i|, \forall i$, $\eta \in \mathcal{S}$. Since T_η is continuous, for all k we find m so that

$$\sup_n \left\{ \sup_{j \geq n} |\theta_{j-n+1}| \frac{b_j^k}{a_n^m} \right\} \leq \sup_n \left\{ \sup_{j \geq n} |\eta_{j-n+1}| \frac{b_j^k}{a_n^m} \right\} < \infty$$

i.e. T_θ is continuous.

Therefore $\theta \in \mathcal{S}$. Hence we obtain that $\mathcal{S}$ is a normal sequence space.

References

[1] L. Crone and W. Robinson, *Diagonal maps and diameters in Köthe spaces*, Israel J. Math. **20**, (1975), 13–22.

[2] G. Köthe, *Topological Vector Spaces 1*, Springer-Verlag, Berlin, Heidelberg, New York, 1969.

[3] R. Meise and D. Vogt, *Introduction to Functional Analysis*, Clarendon Press, Oxford, 1997.

[4] A. Pietsch, *Nuclear Locally Convex Spaces*, Springer-Verlag, Berlin-New York, 1972.

[5] M. S. Ramanujan and T. Terzioğlu, *Subspaces of smooth sequence spaces*, Studia Math. **65**, (1979), 299–312.

[6] T. Terzioğlu, *Die diametrale Dimension von lokalkonvexen Räumen*, Collect. Math. **20**, (1969), 49–99.

Elif Uyanik, Department of Mathematics, Middle East Technical University, 06800 Ankara Turkey

E-mail address: euyanik@metu.edu.tr

Murat H. Yurdakul, Department of Mathematics, Middle East Technical University, 06800 Ankara Turkey

E-mail address: myur@metu.edu.tr